

Life Stage: Prenatal	Target: Mother Reported by: Mother
Administration: Questionnaire <input checked="" type="checkbox"/> Self-administered <input checked="" type="checkbox"/> Researcher/Interviewer-administered	Translation: English
Protocol Measurement Type: Recommended, Preferred	

Brief Explanation of Tool:

The Pregnancy-Unique Quantification of Emesis (PUQE) questionnaire is used to assess the severity of nausea and vomiting in pregnancy (NVP) based on three physical symptoms: nausea, vomiting, and retching over the previous 24 hours.

Equipment and Supplies Needed:

- Questionnaire
 - Paper and pencil for paper administration
 - Electronic data collection platform where this measure is available (Ex: REDCap)

Time to Complete: 2 min

Procedures:

STUDY STAFF INSTRUCTION: This form is recommended to be completed by the pregnant woman during the prenatal life stage using the woman's pregnancy ID.

Please refer to best practices for questionnaire administration. Form completion instructions embedded on form.

References:

Adapted from the 24-hour Pregnancy-Unique Quantification of Emesis (PUQE-24) Scale. Ebrahimi N, Maltepe C, Bournissen FG, Koren G. Nausea and vomiting of pregnancy: Journal of Obstetrics and Gynaecology. 2009;31(9):803-807. doi: 10.1016/S1701-2163(16)34298-0 <https://www.ncbi.nlm.nih.gov/pubmed/19941704>

Location:

<https://echoportal.org/>